

## Author index

- Alejo-Blanco, R., see Kawamoto, F., 101  
 Alexander, K., Parail, A.C. and Parsons, M.  
   An allele of *Trypanosoma brucei* cytoplasmic phosphoglycerate kinase is a mosaic of other alleles and genes (*Short Communication*), 293  
 Aline Jr., R.F., see Smiley, B.L., 143  
 Arnold, H.H., see Ragge, K., 93  
 Åslund, L., see Henriksson, J., 213  
 Avila, H., Goncalves, A.M., Nehme, N.S., Morel, C.M. and Simpson, L.  
   Schizodeme analysis of *Trypanosoma cruzi* stocks from South and Central America by analysis of polymerase chain reaction-amplified minicircle variable region sequences, 175  
 Baack, B.R., see Vander Jagt, D.L., 277  
 Barrett, J., see Brophy, P.M., 205  
 Battistutta, D., see Saul, A., 55  
 Beck, D.J., see Lew, A.M., 153  
 Beverley, S.M. and Coburn, C.M.  
   Recurrent de novo appearance of small linear DNAs in *Leishmania major* and relationship to extrachromosomal DNAs in other species, 133  
 Bobek, L.A., see Henkle, K.J., 69  
 Borst, P., see Crozatier, M., 1  
 Bosch, M., see Gonzalez Clemente, M. del C., 289  
 Bossio, J.C., see Gonzalez Clemente, M. del C., 289  
 Brodin, T.N., see Sacks, D.L., 225  
 Brophy, P.M. and Barrett, J.  
   Strategies for detoxification of aldehydic products of lipid peroxidation in helminths, 205  
 Bzik, D.J., see Tanaka, M., 83  
 Cain, G.D., see Henkle, K.J., 69  
 Campos, N.M., see Vander Jagt, D.L., 277  
 Carr, D., see Thomas, A.W., 285  
 Cazzulo, J.J., see Henriksson, J., 213  
 Coburn, C.M., see Beverley, S.M., 133  
 Conrad, J.T., see Nash, T.E., 125  
 Cook, G.A., see Henkle, K.J., 69  
 Coombs, G.H., see Robertson, C.D., 269  
 Cowman, A.F. and Lew, A.M.  
   Chromosomal rearrangements and point mutations in the DHFR-TS gene of *Plasmodium chabaudi* under antifolate selection, 21  
 Crozatier, M., Van der Ploeg, L.H.T., Johnson, P.J., Gommers-Ampt, J. and Borst, P.  
   Structure of a telomeric expression site for variant specific surface antigens in *Trypanosoma brucei*, 1  
 Dietrich, P., Pilar Dussan, M. del, Floeter-Winter, L.M., Trebilcock Affonso, M.H., Plessman Camargo, E. and Soares, M.B.  
   Restriction fragment length polymorphisms in the ribosomal gene spacers of *Trypanosoma cruzi* and *Trypanosoma conorhini*, 13  
 Docampo, R., see Vercesi, A.E., 119  
 Donelson, J.E., see Henkle, K.J., 69  
 Edwards, M.R., see Grote, R., 109  
 Engman, D.M., see Henkle, K.J., 69  
 Ferrero, M.A.G., see Gonzalez Clemente, M. del C., 289  
 Fetterer, R.H., see Rhoads, M.L., 155  
 Fleck, S.L., see Kawamoto, F., 101  
 Floeter-Winter, L.M., see Dietrich, P., 13  
 Foster, L.A., see Henkle, K.J., 69  
 Franke de Cazzulo, B.M., see Henriksson, J., 213  
 Frasch, A.C.C., see Henriksson, J., 213  
 Gadelha, F.R., see Vercesi, A.E., 119  
 Gamble, H.R., see Zarlenga, D.S., 165  
 Garside, L., see Gibson, W., 45  
 Gibson, W. and Garside, L.  
   Kinetoplast DNA minicircles are inherited from both parents in genetic hybrids of *Trypanosoma brucei*, 45  
 Gommers-Ampt, J., see Crozatier, M., 1  
 Goncalves, A.M., see Avila, H., 175  
 Gonzalez Clemente, M. del C., Ferrero, M.A.G., Bossio, J.C., Bosch, M. and Parodi, A.J.  
   The structure of a galactofuranose-containing oligosaccharide isolated from glycoproteins of the trypanosomatid *Herpetomonas samuelpessoai* (*Short Communication*), 289  
 Gregory, W.F., see Selkirk, M.E., 31  
 Grote, R., Edwards, M.R., Norton, R.S. and O'Sullivan, W.J. A <sup>31</sup>P nuclear magnetic resonance study of *Crithidia luciliae*, 109  
 Gu, H.M., see Tanaka, M., 83  
 Henkle, K.J., Cook, G.A., Foster, L.A., Engman, D.M., Bobek, L.A., Cain, G.D. and Donelson, J.E.  
   The gene family encoding eggshell proteins of *Schistosoma japonicum*, 69  
 Henriksson, J., Åslund, L., Macina, R.A., Franke de Cazzulo, B.M., Cazzulo, J.J., Frasch, A.C.C. and Pettersson, U.  
   Chromosomal localisation of seven cloned antigen genes provides evidence of diploidy and further demonstration of karyotype variability in *Trypanosoma cruzi*, 213  
 Herwaldt, B.L., Schlesinger, P.H. and Krogstad, D.J.  
   Accumulation of chloroquine by membrane preparations from *Plasmodium falciparum*, 257  
 Holder, A., see Keen, J., 241  
 Howard, R.F.  
   The lower-molecular-weight protein complex (R<sub>1</sub>) of the *Plasmodium falciparum* rhoptries lacks the glycolytic enzyme aldolase, 235  
 Hundt, E., see Ragge, K., 93  
 Hunsaker, L.A., see Vander Jagt, D.L., 277  
 Inselburg, J., see Tanaka, M., 83

- Jenkins, R.E., see Selkirk, M.E., 31  
 Jensen, J., see Petersen, C., 187  
 Johnson, P.J., see Crozatier, M., 1
- Kawamoto, F., Alejo-Blanco, R., Fleck, S.L., Kawamoto, Y. and Sinden, R.E.  
 Possible roles of  $\text{Ca}^{2+}$  and cGMP as mediators of the exflagellation of *Plasmodium berghei* and *Plasmodium falciparum*, 101  
 Kawamoto, Y., see Kawamoto, F., 101  
 Keen, J., Holder, A., Playfair, J., Lockyer, M. and Lewis, A.  
 Identification of the gene for a *Plasmodium yoelii* rhoptry protein. Multiple copies in the parasite genome, 241  
 Knapp, B., see Ragge, K., 93  
 Krogstad, D.J., see Herwaldt, B.L., 257
- Leech, J., see Petersen, C., 187  
 Lew, A.M., see Cowman, A.F., 21  
 Lew, A.M. and Beck, D.J.  
 The epitope of a protective monoclonal antibody occurs in a region of microheterogeneity in *Plasmodium chabaudi* (Short Communication), 153  
 Lewis, A., see Keen, J., 239  
 Li, W.-B., see Tanaka, M., 83  
 Lima, S.A., see Vercesi, A.E., 119  
 Lingelbach, K., see Ragge, K., 93  
 Lloyd, D., see Paget, T.A., 63  
 Lockyer, M., see Keen, J., 241
- Macedo, D.V., see Vercesi, A.E., 119  
 Macina, R.A., see Henriksson, J., 213  
 Maizels, R.M., see Selkirk, M.E., 31  
 Merritt, Jr., J.W., see Nash, T.E., 125  
 Morel, C.M., see Avila, H., 175  
 Myler, P.J., see Smiley, B.L., 143
- Nash, T.E., Conrad, J.T. and Merritt Jr., J.W.  
 Variant specific epitopes of *Giardia lamblia*, 125  
 Nehme, N.S., see Avila, H., 175  
 Nelson, R., see Petersen, C., 189  
 Nielsen, B., see Parsons, M., 197  
 Norton, R.S., see Grote, R., 109
- O'Sullivan, W.J., see Grote, R., 109
- Paget, T.A., Raynor, M.H., Shipp, D.W.E. and Lloyd, D.  
*Giardia lamblia* produces alanine anaerobically but not in the presence of oxygen, 63  
 Parail, A.C., see Alexander, K., 293  
 Parodi, A.J., see Gonzalez Clemente, M. del C., 289  
 Parsons, M. and Nielsen, B.  
 Active transport of 2-deoxy-D-glucose in *Trypanosoma brucei* procyclic forms, 197  
 Parsons, M., see Alexander, K., 293  
 Pedraza, G., see Shapira, M., 247  
 Petersen, C., Nelson, R., Leech, J., Jensen, J., Wollish, W. and Scherf, A.  
 The gene product of the *Plasmodium falciparum* 11.1 locus is a protein larger than one megadalton, 189  
 Pettersson, U., see Henriksson, J., 213  
 Pilar Dussan, M. del, see Dietrich, P., 13  
 Playfair, J., see Keen, J., 241  
 Plessman Camargo, E., see Dietrich, P., 13
- Ragge, K., Arnold, H.-H., Tummler, M., Knapp, B., Hundt, E. and Lingelbach, K.  
 In vitro biosynthesis and membrane translocation of the serine rich protein of *Plasmodium falciparum*, 93  
 Raynor, M.H., see Paget, T.A., 63  
 Rhoads, M.L. and Fetterer, R.H.  
 Biochemical and immunochemical characterization of  $^{125}\text{I}$ -labeled cuticle components of *Haemonchus contortus*, 155  
 Robertson, C.D. and Coombs, G.H.  
 Characterisation of three groups of cysteine proteinase in the amastigotes of *Leishmania mexicana mexicana*, 269
- Sacks, D.L., Brodin, T.N. and Turco, S.J.  
 Developmental modification of the lipophosphoglycan from *Leishmania major* promastigotes during metacyclogenesis, 225  
 Saul, A. and Battistutta, D. Analysis of the sequences flanking the translational start sites of *Plasmodium falciparum*, 55  
 Scherf, A., see Petersen, C., 189  
 Schlesinger, P.H., see Herwaldt, B.L., 257  
 Selkirk, M.E., Gregory, W.F., Yazdanbakhsh, M., Jenkins, R.E. and Maizels, R.M.  
 Cuticular localisation and turnover of the major surface glycoprotein (gp29) of adult *Brugia malayi*, 31  
 Shapira, M. and Pedraza, G.  
 Sequence analysis and transcriptional activation of heat shock protein 83 of *Leishmania mexicana amazonensis*, 247  
 Shipp, D.W.E., see Paget, T.A., 63  
 Simpson, L., see Avila, H., 175  
 Sinden, R.E., see Kawamoto, F., 101  
 Smiley, B.L., Aline, Jr., R.F., Myler, P.J. and Stuart, K.  
 A retroposon in the 5' flank of a *Trypanosoma brucei* VSG gene lacks insertional terminal repeats, 143  
 Soares, M.B., see Dietrich, P., 13  
 Stuart, K., see Smiley, B.L., 143
- Tanaka, M., Gu, H.-M., Bzik, D.J., Li, W.-B. and Inselburg, J.  
 Mutant dihydrofolate reductase-thymidylate synthase genes in pyrimethamine-resistant *Plasmodium falciparum* with polymorphic chromosome duplications, 83  
 Thomas, A.W., Waters, A.P. and Carr, D.  
 Analysis of variation in PF83, an erythrocytic merozoite vaccine candidate antigen of *Plasmodium falciparum* (Short Communication), 285  
 Trebilcock Affonso, M.H., see Dietrich, P., 13  
 Tummler, M., see Ragge, K., 93  
 Turco, S.J., see Sacks, D.L., 225
- Van der Ploeg, L.H.T., see Crozatier, M., 1  
 Vander Jagt, D.L., Hunsaker, L.A., Campos, N.M. and Baack, B.R.  
 D-Lactate production in erythrocytes infected with *Plasmodium falciparum*, 277  
 Vercesi, A.E., Macedo, D.V., Lima, S.A., Gadelha, F.R. and Docampo, R.  
 $\text{Ca}^{2+}$  transport in digitonin-permeabilized trypanosomatids, 119
- Waters, A.P., see Thomas, A.W., 285  
 Wollish, W., see Petersen, C., 189
- Yazdanbakhsh, M., see Selkirk, M.E., 31
- Zarlenga, D.S. and Gamble, H.R.  
 Molecular cloning and expression of an immunodominant 53-kDa excretory-secretory antigen from *Trichinella spiralis* muscle larvae, 165

## Subject index

- Acidification, 257  
 Alcohol dehydrogenase, 205  
 Aldehyde reductase, 205  
 Aldolase, 235  
 Alpha-amanitin-resistant transcription, 1  
 Anaerobic-aerobic transition, 63  
 Antifolate, 21  
 Antigenic variation, 125  
 Antigenic variation, 143  
 Antigenic variation, 285  
  
*Brugia malayi*, 31  
  
 Ca<sup>2+</sup> mobilization, 101  
 Ca<sup>2+</sup> transport, 119  
 cGMP, 101  
 Chloroquine accumulation, 257  
 Chloroquine, 257  
 Chromosomal alteration, 133  
 Chromosome, 21  
 Chromosome size polymorphism, 83  
 Collagen, 155  
*Crithidia fasciculata*, 119  
*Crithidia luciliae*, 109  
 Cuticle, 155  
 Cuticle, 31  
 Cysteine-rich proteins, 125  
 Cytoadherence, 189  
  
 2-Deoxyglucose, 197  
 Dihydrofolate reductase-thymidylate synthase point mutation, 83  
 DNA sequence, 285, 296  
 Drug resistance, 133  
 Drug resistance, 21  
  
 Eggshell protein, 69  
 ES antigen, 163  
 Exflagellation, 101  
 Evolution, 293  
  
 Fermentation product, 63  
  
 Gametogenesis, 101  
 Gene, 241  
 Gene amplification, 133  
 Gene cloning, 163  
 Gene conversion, 293  
 Gene family, 69  
 Genetic exchange, 45  
*Giardia lamblia*, 63  
*Giardia*, 125  
 Glucose metabolism, 109  
 Glucose transporter, 197  
 Glutathione transferase, 205  
  
 Glycoprotein, 289  
 Glycosome, 293  
 Gp29, 31  
  
*Haemonchus contortus*, 155  
 Heat shock protein 83, 247  
 Helminth, 203  
*Herpetomonas samuelpessoai*, 289  
  
 Immunity, 241  
 Ingi, 143  
 Initiation codon, 55  
 Intergenic sequence, 247  
 Intracellular transport, 93  
  
 Kinetics, 197  
 Kinetoplast DNA, 45  
 Kinetoplastida, 109  
  
 D-Lactate, 277  
 L-Lactate, 277  
*Leishmania agamae*, 119  
*Leishmania mexicana amazonensis*, 247  
*Leishmania mexicana*, 119  
*Leishmania mexicana*, 269  
*Leishmania*, 225  
 Lipid peroxidation, 205  
 Lipophosphoglycan, 225  
  
 Malaria, 21, 189, 241, 257  
 Membrane vesicle, 257  
 Merozoite antigen, 285  
 Metacyclogenesis, 225  
 Methotrexate, 133  
 Methylglyoxal, 277  
 Microbody, 293  
 Minicircle DNA, 175  
 Mitochondrial DNA, 45  
 Mitochondrion, 119  
 Molecular karyotype, 133  
 Molecular karyotype, 213  
  
 Nematode, 31  
 Nuclear extract, 69  
  
 Oligosaccharide, 289  
  
<sup>31</sup>P magnetic resonance spectroscopy, 109  
 Parasite diagnosis, 165  
 Peptidyl aminomethylcoumarin, 269  
 PfEMP 1, 189  
 pH, 109  
 Phosphoglycerate kinase, 293  
*Plasmodium berghei*, 101  
*Plasmodium falciparum*, 55, 83, 93, 101, 189, 235, 257, 277, 285



- Plasmodium yoelii*, 241  
Ploidy, 213  
Polymerase chain reaction, 175  
Polyphosphate, 109  
Procyclic form, 197  
Promoter, 1  
Proteinase, 269  
Proton NMR, 63  
Pulsed field gel electrophoresis, 213  
Purification, 269  
Pyrimethamine resistance, 83  
  
Radioiodination, 155  
Recombination, 143  
Restriction fragment length polymorphism, 13  
Retroposon, 143  
Rhoptry, 235  
Rhoptry, 241  
R<sub>I</sub> protein complex, 235  
Ring-infected erythrocyte surface antigen, 189  
  
Schistosome, 69  
Schizodeme, 175  
  
Secretion, 93  
Serine-rich protein, 93  
Specificity, 269  
Surface antigen, 1, 31, 125  
Swine, 165  
  
Taxonomy, 13  
Transcription, 69  
Translation, 55  
Trichinellosis, 165  
Trypanosomatid, 289  
*Trypanosoma brucei*, 45, 197, 293  
*Trypanosoma conorhini* rRNA genes, 13  
*Trypanosoma cruzi* rRNA genes, 13  
*Trypanosoma cruzi*, 175, 213  
Trypanosome repeated sequence, 143  
Trypanosome, 1  
  
UV light transcription mapping, 1  
  
Vaccine, 285

